

AD_____

Award Number: W81XWH-12-1-0605

TITLE: Molecular Indicators of Castration-resistant Prostate Cancer

PRINCIPAL INVESTIGATOR: Jun Luo, Ph.D.

CONTRACTING ORGANIZATION: Johns Hopkins University,
Baltimore, MD 21218-2680

REPORT DATE: October 2013

TYPE OF REPORT: Annual report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE October 06, 2013		2. REPORT TYPE Annual		3. DATES COVERED Sept. 30 th , 2012 – Sept. 29 th , 2013	
4. TITLE AND SUBTITLE Molecular indicators of castration-resistant prostate cancer				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-12-1-0605	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jun Luo, Ph.D. Betty Diamond E-Mail: jluo1@jhmi.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Johns Hopkins University Baltimore, MD 21218-2680				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Metastatic prostate cancers are commonly treated by agents designed to suppress androgen receptor (AR) signaling mediated by the full-length AR (AR-FL). Why some patients progress rapidly after treatment while others benefit with prolonged remission is an unsolved question. We propose approaches to develop molecular indicators of response and resistance that will enable prediction (before therapy) or early detection (during therapy) of therapeutic benefit. We will test the hypothesis that AR splice variants (AR-Vs) are molecular indicators of castration-resistant prostate cancer (CRPC). During this funding period, we achieved a major milestone by completing experimental phase of the proposed transcriptome sequencing studies, and began the data analysis phase of the project. In addition, we have identified and acquired biopsy specimens that will be required for the proposed in-situ detection methods. We conclude that androgen receptor splice variants can be detected by RNA sequencing in clinical specimens derived from patients with castration-resistant prostate cancer.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	2
Reportable Outcomes.....	2
Conclusion.....	3
References.....	3
Appendices.....	4
Supporting Data.....	4

Introduction

Hormone therapies block androgen production and/or androgen receptor (AR) function leading to a period of clinical regression varying from months to more than 6 years among patients treated for metastatic prostate cancer. Because restored AR signaling is the key determinant of castration-resistant prostate cancer (CRPC), aberrant variants of the androgen receptor may mediate and indicate therapeutic resistance. Our recent efforts (1-3) have established that truncated androgen receptor splice variants (AR-Vs) are resistant to castration therapies that target the canonical full-length AR (AR-FL). We are now equipped with the knowledge that AR-Vs are both structurally and functionally distinguishable from the full-length AR (AR-FL), which remains the key target in prostate cancer drug design. We view these new discoveries and knowledge as opportunities that have yet to be capitalized to address the unmet need of developing indicators of castration resistance. In our previous studies, we have shown elevated expression of the AR-Vs following suppression of AR-FL (3). The primary purpose of the project is to discover and develop molecular indicators of CRPC by targeting the AR-Vs. We will test the hypothesis that AR-Vs are molecular indicators of CRPC. The scope of the proposed research is: 1) to perform RNA-Seq targeting transcripts originated from the human AR gene locus for *ab initio* cataloging of AR-Vs that define CRPC; and 2) to establish the proof-of-principle that AR-Vs detected prior to the initiation of hormone therapies may be used to predict CRPC progression.

Body

Findings resulting from Task 1: To employ a novel targeted RNA-Seq method for *ab initio* cataloging of androgen receptor variants that define CRPC (Months 1-24).

Summary: By generating raw sequencing data using RNA-seq from 48 clinical specimens, we have achieved a critical project milestones associated with Task 1 as outlined in SOW. As stated in SOW, data analysis involving all specimens will be performed during the second year of the funding period. On the basis of initial findings from three specimens, we conclude that AR-V7, one of the most important androgen receptor splice variant (4), is abundantly expressed in CRPC and detected by RNA-seq.

Supporting data: Quantitative RT-PCR results suggest that the expression level of each individual AR-V is often a small fraction of the total level for AR-FL. For example, AR-V7 represents only about 1% to 5% of the total AR in the vast majority of CRPC specimens examined (1, 5). Because of the lower abundance of AR-V than AR-FL, the clinical significance and functional role of AR-V has been debated (4-6). However, we posit that 1) accurate quantification of AR-V transcripts would be compromised by unknown splice junctions as well as various technical constraints of the conventional RT-PCR methodology and 2) even at a small fraction of the AR-FL, AR-V expression in treatment-resistant specimens may reach levels equivalent to those of AR-FL in untreated tumors because AR-FL is often massively overexpressed in CRPC (7), thus conferring

functional significance when AR-FL is suppressed. To address accurate quantification, we performed RNA-seq in three independent CRPC specimens using the Illumina Hi-Seq 2000 platform (Figure 1). Comparing to qRT-PCR data from the same specimens, we found that RT-PCR *underestimated* the AR-V7 abundance by an average of 5 fold, and that in one CRPC specimen the ratio of AR-V7 to AR-FL reached 41%, using the number of AR-FL- and AR-V7-specific exon-spanning sequence reads as proxies for relative transcript abundance (Figure 1). Taken together, these data suggest that AR-V7 is a high-abundance transcript in a subset of clinical CRPC, and may be readily detected in CRPC cells. Other known AR-Vs are either orders of magnitude lower in abundance than AR-V7, or functionally inactive (2). Another constitutively active AR-V, ARV567ES, is frequently expressed in CRPC (8). Interestingly, splice junctions specific for ARV567ES are not detected by the three RNA samples sequenced so far (Figure 1). These findings suggest that although other variants may contribute to CRPC, AR-V7 is the best candidate for designing detection assays to be used in clinical specimens.

Findings resulting from Task 2: To establish the proof-of-principle that AR-Vs detected prior to the initiation of hormone therapies may be used to predict CRPC progression (Months 1-36).

According to our project plan in SOW we will focus on sample preparation during year 1 of the project. Experimental data and findings will not be available until later years of the project period.

Key Research Accomplishments

1. Completed raw RNA-seq data collection from 48 samples.
2. Established that RNA-seq can be used to reliably estimate the abundance of AR variant transcripts.

Reportable Outcomes

Manuscripts: None at this time.

Presentations: None at this time.

Grant Applications:

Title: The aberrant androgen receptor underlies abiraterone/enzalutamide resistance

Supporting Agency: NIH (Luo - PI)

Name of Procuring Contracting/Grants Officer: TBD - NIH

Address of Funding Agency: 9000 Rockville Pike, Bethesda, MD 20892

Performance Period: 4/1/2014 - 3/31/2019

Status: Submitted on June 5th, 2013

Conclusion

We conclude that androgen receptor splice variants can be detected by RNA sequencing in clinical specimens derived from patients with castration-resistant prostate cancer. Our findings suggest that previous methods of detection may have *underestimated* the AR splice variant transcript abundance. The implication is that AR splice variants may be more abundant than previously thought, allowing robust detection in clinical specimens.

References

1. Hu R, Dunn TA, Wei S, Isharwal S, Veltri RW, Humphreys E, Han M, Partin AW, Vessella RL, Isaacs WB, Bova GS, Luo J. Ligand-independent androgen receptor variants derived from splicing of cryptic exons signify hormone-refractory prostate cancer. *Cancer Res.* 2009;69(1):16-22. PMCID: 2614301.
2. Hu R, Isaacs WB, Luo J. A snapshot of the expression signature of androgen receptor splicing variants and their distinctive transcriptional activities. *Prostate.* 2011;71(15):1656-67. PMCID: 3360954.
3. Hu R, Lu C, Mostaghel EA, Yegnasubramanian S, Gurel M, Tannahill C, Edwards J, Isaacs W, Nelson PS, Bluemn E, Plymate SR, Luo J. Distinct transcriptional programs mediated by the ligand-dependent full-length androgen receptor and its splice variants in castration-resistant prostate cancer. *Cancer Res.* 2012;accepted.
4. Plymate SR, Luo J. The Expression Signature of Androgen Receptor Splice Variants and Their Distinctive Transcriptional Activities in Castration-Resistant Prostate Cancer. *Androgen-Responsive Genes in Prostate Cancer*: Springer; 2013. p. 201-13.
5. Watson PA, Chen YF, Balbas MD, Wongvipat J, Socci ND, Viale A, Kim K, Sawyers CL. Constitutively active androgen receptor splice variants expressed in castration-resistant prostate cancer require full-length androgen receptor. *Proc Natl Acad Sci U S A.* 2010;107(39):16759-65. PMCID: 2947883.
6. Sadar MD. Small molecule inhibitors targeting the "achilles' heel" of androgen receptor activity. *Cancer Res.* 2011;71(4):1208-13. PMCID: 3132148.
7. Isaacs JT, D'Antonio JM, Chen S, Antony L, Dalrymple SP, Ndikuyeze GH, Luo J, Denmeade SR. Adaptive auto-regulation of androgen receptor provides a paradigm shifting rationale for bipolar androgen therapy (BAT) for castrate resistant human prostate cancer. *Prostate.* 2012;72(14):1491-505. PMCID: 3374010.
8. Sun S, Sprenger CC, Vessella RL, Haugk K, Soriano K, Mostaghel EA, Page ST, Coleman IM, Nguyen HM, Sun H, Nelson PS, Plymate SR. Castration resistance in human prostate cancer is conferred by a frequently occurring androgen receptor splice variant. *J Clin Invest.* 2010;120(8):2715-30. PMCID: 2912187.

Appendices

None

Supporting Data (Figure 1 and legend)

Figure 1. RNA-seq results illustrating the splice junction tracks for the androgen receptor locus in 3 CRPC specimens. The number of reads specific to each splice junction were compared with data derived from conventional RT-PCR. For example, the number of reads spanning AR-V7 splice junction (second blue arrow) is 41 in CRPC #2 and reads spanning AR-FL splice junction is 111, while the V7/FL ratio in this sample by QRT-PCR is 8%, suggesting QRT-PCR underestimates the true ratio of AR-V7/AR-FL by 5 fold. Blue arrows indicate splice junctions. The number of reads were extracted from 3 specimens (CRPC #1, #2, and #3).

